



# EX NAVODAYAN FOUNDATION

(Registered Under Indian Trust Act 1882)

Reg. No. : 2016, 43

B/36/43 46M Brahmanand Colony, Durgakund, Varanasi (UP) 221005

Mob.: 8958143003

Email Id : exnavodayanfoundation@gmail.com

30/01/2023

Morning

## Answers & Solutions

Time : 3 hrs.

*for*

M.M. : 300

## JEE (Main)-2023 (Online) Phase-1

(Physics, Chemistry and Mathematics)

### IMPORTANT INSTRUCTIONS:

- (1) The test is of **3 hours** duration.
- (2) The Test Booklet consists of 90 questions. The maximum marks are 300.
- (3) There are **three** parts in the question paper consisting of **Physics, Chemistry** and **Mathematics** having 30 questions in each part of equal weightage. Each part (subject) has two sections.
  - (i) **Section-A:** This section contains 20 multiple choice questions which have only one correct answer. Each question carries **4 marks** for correct answer and **-1 mark** for wrong answer.
  - (ii) **Section-B:** This section contains 10 questions. In Section-B, attempt any **five questions out of 10**. The answer to each of the questions is a numerical value. Each question carries **4 marks** for correct answer and **-1 mark** for wrong answer. For Section-B, the answer should be rounded off to the nearest integer.

## PHYSICS

### SECTION - A

**Multiple Choice Questions:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

**Choose the correct answer:**

1. Choose the correct relationship between Poisson ratio ( $\sigma$ ), bulk modulus ( $K$ ) and modulus of rigidity ( $\eta$ ) of a given solid object

(1)  $\sigma = \frac{6K - 2\eta}{3K - 2\eta}$       (2)  $\sigma = \frac{6K + 2\eta}{3K - 2\eta}$

(3)  $\sigma = \frac{3K - 2\eta}{6K + 2\eta}$       (4)  $\sigma = \frac{3K + 2\eta}{6K + 2\eta}$

**Answer (3)**

**Sol.** Poisson ratio ( $\sigma$ ), bulk modulus ( $K$ ) and modulus of rigidity ( $\eta$ ) are related by

$$\therefore 2\eta(1 + \sigma) = 3K(1 - 2\sigma)$$

$$2\eta + 2\eta\sigma = 3K - 6K\sigma$$

$$\sigma = \frac{3K - 2\eta}{2\eta + 6K}$$

2. A small object at rest, absorbs a light pulse of power 20 mW and duration 300 ns. Assuming speed of light as  $3 \times 10^8$  m/s, the momentum of the object becomes equal to


(1)  $2 \times 10^{-17}$  kg m/s      (2)  $3 \times 10^{-17}$  kg m/s  
 (3)  $1 \times 10^{-17}$  kg m/s      (4)  $0.5 \times 10^{-17}$  kg m/s

**Answer (1)**

**Sol.** Assuming the small object as photon.

$$\begin{aligned} \text{Momentum } (p) &= \frac{E}{C} \\ &= \frac{20 \times 10^{-3} \times 300 \times 10^{-9}}{3 \times 10^8} \\ &= 2 \times 10^{-17} \text{ kg m/s} \end{aligned}$$

3. A massless square loop, of wire of resistance  $10 \Omega$ , supporting a mass of 1 g, hangs vertically with one of its sides in a uniform magnetic field of  $10^3$  G, directed outwards in the shaded region. A dc voltage  $V$  is applied to the loop. For what value of  $V$ , the magnetic force will exactly balance the weight of the supporting mass of 1 g? (If sides of the loop = 10 cm,  $g = 10$  m/s<sup>2</sup>)



- (1) 1 V      (2) 10 V  
 (3)  $\frac{1}{10}$  V      (4) 100 V

**Answer (2)**

**Sol.** For balancing of force

$$\therefore F_{\text{loop}} = \text{weight}$$

$$\left(\frac{V}{R}\right)lB = mg$$

$$\left(\frac{V}{10}\right) \times \frac{10}{100} \times (10^3 \times 10^{-4}) = \left(\frac{1}{1000}\right) \times 10$$

$$V = 10 \text{ volts}$$

4. The magnetic moment associated with two closely wound circular coils  $A$  and  $B$  of radius  $r_A = 10$  cm and  $r_B = 20$  cm respectively are equal if: (where  $N_A$ ,  $I_A$  and  $N_B$ ,  $I_B$  are number of turn and current of  $A$  and  $B$  respectively)

- (1)  $N_A I_A = 4 N_B I_B$       (2)  $2 N_A I_A = N_B I_B$   
 (3)  $N_A = 2 N_B$       (4)  $4 N_A I_A = N_B I_B$

**Answer (1)**

**Sol.**  $M_A = M_B$

$$I_A N_A (\pi r_A^2) = I_B N_B (\pi r_B^2)$$


$$I_A N_A = 4 I_B N_B$$

5. A ball of mass 200 g rests on a vertical post of height 20 m. A bullet of mass 10 g, travelling in horizontal direction, hits the centre of the ball. After collision both travels independently. The ball hits the ground at a distance 30 m and bullet at a distance of 120 m from the foot of the post. The value of initial velocity of the bullet will be (if  $g = 10$  m/s<sup>2</sup>)

- (1) 60 m/s      (2) 120 m/s  
 (3) 400 m/s      (4) 360 m/s

**Answer (4)**






Choose the correct answer from the options given below :


- (1) A-I, B-III, C-IV, D-II
- (2) A-II, B-III, C-IV, D-I
- (3) A-I, B-II, C-III, D-IV
- (4) A-II, B-IV, C-III, D-I

**Answer (4)**

Sol.




10. The output waveform of the given logical circuit for the following inputs A and B as shown below, is



**Answer (1)**


Sol.



Truth table

A	B	Y
0	0	0
0	1	1
1	0	1
1	1	1

OR Gate



11. The charge flowing in a conductor changes with time as  $Q(t) = \alpha t - \beta t^2 + \gamma t^3$ . Where  $\alpha$ ,  $\beta$  and  $\gamma$  are constants. Minimum value of current is

- (1)  $\alpha - \frac{\gamma^2}{3\beta}$                       (2)  $\alpha - \frac{\beta^2}{3\gamma}$   
 (3)  $\alpha - \frac{3\beta^2}{\gamma}$                       (4)  $\beta - \frac{\alpha^2}{3\gamma}$

**Answer (2)**

**Sol.**  $Q(t) = \alpha t - \beta t^2 + \gamma t^3$

$$i(t) = \alpha - 2\beta t + 3\gamma t^2$$

$$\frac{di}{dt} = -2\beta + 6\gamma t = 0 \quad (\text{for max/min of } i)$$

at  $t = \frac{\beta}{3\gamma}$  ( $i$  is minimum as  $i$  is an upward parabola)

$$i\left(\frac{\beta}{3\gamma}\right) = \alpha - 2\beta\left(\frac{\beta}{3\gamma}\right) + \frac{3\gamma\beta^2}{9\gamma^2}$$

$$= \alpha - \frac{\beta^2}{3\gamma}$$

12. The height of liquid column raised in a capillary tube of certain radius when dipped in liquid A vertically is, 5 cm. If the tube is dipped in a similar manner in another liquid B of surface tension and density double the values of liquid A, the height of liquid column raised in liquid B would be \_\_\_\_\_ m.

- (1) 0.20                      (2) 0.05  
 (3) 0.5                      (4) 0.10

**Answer (2)**

**Sol.** height of capillary rise =  $\frac{2s \cos \theta}{\rho g R}$

$$\text{When in A } 5 \text{ cm} = \frac{2s_A \cos \theta}{\rho_A g R}$$

$$\text{When in B } h = \frac{2s_B \cos \theta}{\rho_B g R}$$

$$s_B = 2s_A \text{ and } \rho_B = 2\rho_A$$

$$h = \frac{2 \times 2s_A \times \cos \theta}{2\rho_A g R} = 5 \text{ cm}$$

13. A person has been using spectacles of power  $-1.0$  dioptre for distant vision and a separate reading glass of power 2.0 dioptres. What is the least distance of distinct vision for this person

- (1) 50 cm                      (2) 10 cm  
 (3) 30 cm                      (4) 40 cm

**Answer (1)**

**Sol.**  $u = 25 \text{ cm}$

$$f = \frac{1}{2} \text{ m} = 50 \text{ cm}$$

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

$$\Rightarrow \frac{1}{v} + \frac{1}{25} = -\frac{1}{50}$$

$$\frac{1}{v} = -\frac{1}{50}$$

$$\Rightarrow u = -50 \text{ cm}$$

14. Speed of an electron in Bohr's 7<sup>th</sup> orbit for Hydrogen atom is  $3.6 \times 10^6 \text{ m/s}$ . The corresponding speed of the electron in 3<sup>rd</sup> orbit, in m/s is

- (1)  $(7.5 \times 10^6)$                       (2)  $(1.08 \times 10^6)$   
 (3)  $(8.4 \times 10^6)$                       (4)  $(3.6 \times 10^6)$

**Answer (3)**

**Sol.**  $v \propto \frac{Z}{n}$

$$\frac{v_1}{v_2} = \left(\frac{n_2}{n_1}\right)$$

$$\Rightarrow \frac{3.6 \times 10^6}{v_2} = \frac{3}{7}$$

$$\Rightarrow v_2 = \frac{7}{3} \times 3.6 \times 10^6 \text{ m/s}$$

$$= 8.4 \times 10^6 \text{ m/s}$$

15. Electric field in a certain region is given by

$$\vec{E} = \left(\frac{A}{x^2} \hat{i} + \frac{B}{y^2} \hat{j}\right). \text{ The SI unit of A and B are}$$

- (1)  $\text{Nm}^3\text{C}; \text{Nm}^2\text{C}$   
 (2)  $\text{Nm}^2\text{C}; \text{Nm}^3\text{C}$   
 (3)  $\text{Nm}^2\text{C}^{-1}; \text{Nm}^2\text{C}^{-1}$   
 (4)  $\text{Nm}^3\text{C}^{-1}; \text{Nm}^2\text{C}^{-1}$

**Answer (3)**

Sol.  $\vec{E} = \left( \frac{A}{x^2} \hat{i} + \frac{B}{y^3} \hat{j} \right)$

$\left[ \frac{A}{x^2} \right] = [E] = \left[ \frac{F}{q} \right] = \left[ \frac{N}{C} \right] = NC^{-1}$

$[A] = (Nm^2C^{-1})$

$[B] = Nm^3C^{-1}$

16. A sinusoidal carrier voltage is amplitude modulated. The resultant amplitude modulated wave has maximum and minimum amplitude of 120 V and 80 V respectively. The amplitude of each sideband is

- (1) 10 V                                  (2) 15 V  
 (3) 20 V                                  (4) 5 V

**Answer (1)**

Sol. Amplitude of each side band =  $\frac{A_{\text{message}}}{2}$

$A_{\text{carrier}} + A_{\text{message}} = 120$                                   ... (1)


$A_{\text{carrier}} - A_{\text{message}} = 80$                                   ... (2)

From (1) and (2)

$A_{\text{message}} = 20 \text{ V}$


∴ Amplitude of each side band = 10 V

17. As per the given figure, a small ball P slides down the quadrant of a circle and hits the other ball Q of equal mass which is initially at rest. Neglecting the effect of friction and assume the collision to be elastic, the velocity of ball Q after collision will be ( $g = 10 \text{ m/s}^2$ )



- (1) 0.25 m/s                                  (2) 2 m/s  
 (3) 0    (4) 4 m/s

**Answer (2)**



Mass is same and elastic collision, so speed gets exchanged,  $v = 2 \text{ m/s}$

18. If the gravitational field in the space is given as  $\left( -\frac{K}{r^2} \right)$ . Taking the reference point to be at  $r = 2 \text{ cm}$  with gravitational potential  $V = 10 \text{ J/kg}$ . Find the gravitational potential at  $r = 3 \text{ cm}$  in SI unit (Given, that  $K = 6 \text{ Jcm/kg}$ )

- (1) 10  
 (2) 12  
 (3) 11  
 (4) 9

**Answer (3)**

Sol.  $E = -\frac{K}{r^2}$

$\Delta V = -\int_{r=2 \text{ cm}}^{3 \text{ cm}} E \cdot dr$

$= \int_2^3 \frac{k}{r^2} dr$

$= \left[ -\frac{K}{r} \right]_2^3 = \left( \frac{K}{6} \right) = \frac{6}{6} = 1 \text{ J/kg}$

$V_f - V_i = 1$

$\Rightarrow V_f - 10 = 1$

$V_f = 11 \text{ J/kg}$

19. In a series LR circuit with  $X_L = R$ , power factor is  $P_1$ . If a capacitor of capacitance  $C$  with  $X_C = X_L$  is added to the circuit the power factor becomes  $P_2$ . The ratio of  $P_1$  to  $P_2$  will be:

- (1) 1 : 2  
 (2) 1 : 3  
 (3) 1 :  $\sqrt{2}$   
 (4) 1 : 1

**Answer (3)**

Sol.  $X_L = R$

$\Rightarrow P_1 = \frac{R}{\sqrt{X_L^2 + R^2}} = \frac{1}{\sqrt{2}}$


Now,  $X_L = X_C = R$

$\Rightarrow P_2 = \frac{R}{\sqrt{R^2 + (X_L - X_C)^2}} = 1$

$\Rightarrow \frac{P_1}{P_2} = \frac{1}{\sqrt{2}}$

20. The figure represents the momentum time (p-t) curve for a particle moving along an axis under the influence of the force. Identify the regions on the graph where the magnitude of the force is maximum and minimum respectively?

If  $(t_3 - t_2) < t_1$



- (1) a and b                      (2) c and a  
 (3) c and b                      (4) b and c

**Answer (3)**

**Sol.**  $F = \frac{dp}{dt}$

$\Rightarrow |F| = \left| \frac{dp}{dt} \right| = |\text{slope of } p - t \text{ curve}|$


As we can see from graph,

$|F_c|$  is maximum and  $|F_b|$  is minimum.

**SECTION - B**

**Numerical Value Type Questions:** This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g. 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.


21. In Young's double slit experiment, two slits  $S_1$  and  $S_2$  are 'd' distance apart and the separation from slits to screen is D (as shown in figure). Now if two transparent slabs of equal thickness 0.1 mm but refractive index 1.51 and 1.55 are introduced in the path of beam ( $\lambda = 4000 \text{ \AA}$ ) from  $S_1$  and  $S_2$  respectively. The central bright fringe spot will shift by \_\_\_\_\_ number of fringes.



**Answer (10)**

**Sol.** Path difference introduced by two slabs =  $(\mu_2 - \mu_1)t$

$$\begin{aligned} \Rightarrow \text{Number of shifts} &= \frac{(\mu_2 - \mu_1)t}{\lambda} \\ &= \frac{0.04 \times 0.1 \text{ mm}}{4000 \text{ \AA}} \\ &= \frac{4 \times 10^{-2} \times 10^{-4}}{4 \times 10^{-7}} \\ &= 10 \end{aligned}$$



22.

As per the given figure, if  $\frac{di}{dt} = -1 \text{ A/s}$  then the value of  $V_{AB}$  at this instant will be \_\_\_\_\_ V.

**Answer (30)**

**Sol.** From the circuit :

$$V_A - iR - \frac{Ldi}{dt} - 12 = V_B$$

$$\begin{aligned} \Rightarrow V_A - V_B &= 2 \times 12 + 6(-1) + 12 \text{ volts} \\ &= 30 \text{ volts} \end{aligned}$$

23. A horse rider covers half the distance with 5 m/s speed. The remaining part of the distance was travelled with speed 10 m/s for half the time and with speed 15 m/s for other half of the time. The mean speed of the rider averaged over the whole time of motion is  $\frac{x}{7}$  m/s. The value of x is \_\_\_\_\_.

**Answer (50)**

**Sol.** Let S total distance

$$\Rightarrow t_1 = \frac{S}{5} \quad \dots(1)$$

$$\text{Also, } \frac{S}{2} = \frac{10t_2}{2} + \frac{15t_2}{2}$$

$$\Rightarrow t_2 = \frac{S}{25} \quad \dots(2)$$

$$\begin{aligned} \Rightarrow \text{Mean speed} &= \frac{S}{t_1 + t_2} \\ &= \frac{S}{\frac{S}{5} + \frac{S}{25}} = \frac{250}{35} \text{ m/s} = \frac{50}{7} \text{ m/s} \end{aligned}$$

24. In an experiment for estimating the value of focal length of converging mirror, image of an object placed at 40 cm from the pole of the mirror is formed at distance 120 cm from the pole of the mirror. These distances are measured with a modified scale in which there are 20 small divisions in 1 cm. The value of error in measurement of focal length of the mirror is  $\frac{1}{K}$  cm. The value of x is \_\_\_\_\_.

**Answer (32)**

**Sol.**  $\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$  ... (1)

$$\Rightarrow -\frac{1}{f^2} df = -\frac{1}{v^2} dv - \frac{1}{u^2} du$$

$$\Rightarrow \frac{df}{f^2} = \frac{dv}{v^2} + \frac{du}{u^2} \quad \dots(2)$$

From (1):  $-\frac{1}{120} - \frac{1}{40} = \frac{1}{f} \Rightarrow f = -30$  cm

Also, least count =  $\frac{1 \text{ cm}}{20} = 0.05$  cm

$$\Rightarrow df = \left[ \frac{0.05}{120^2} + \frac{0.05}{40^2} \right] \times 30^2$$

$$= 0.05 \left[ \frac{1}{16} + \frac{9}{16} \right] = \frac{5}{8} \times \frac{5}{100} = \frac{1}{32} \text{ cm}$$

$$\Rightarrow k = 32$$

25. In a screw gauge, there are 100 divisions on the circular scale and the main scale moves by 0.5 mm on a complete rotation of the circular scale. The zero of circular scale lies 6 divisions below the line of graduation when two studs are brought in contact with each other. When a wire is placed between the studs, 4 linear scale divisions are clearly visible while 46<sup>th</sup> division the circular scale coincide with the reference line. The diameter of the wire is \_\_\_\_\_  $\times 10^{-2}$  mm.

**Answer (220)**

**Sol.** Least count of screw gauge =  $\frac{0.5}{100}$  mm =  $\frac{1}{200}$  mm

$$\begin{aligned} \text{Zero error of screw gauge} &= +\frac{6}{200} \text{ mm} = +\frac{3}{100} \\ &= 0.03 \text{ mm} \end{aligned}$$

$$\begin{aligned} \text{Reading of screw gauge} &= 4 \times 0.5 + \frac{46}{200} \text{ mm} \\ &= 2 + \frac{23}{100} \text{ mm} = 2.23 \text{ mm} \end{aligned}$$

$$\begin{aligned} \text{So diameter of wire} &= 2.23 \text{ mm} - 0.03 \text{ mm} \\ &= 2.20 \text{ mm} \\ &= 220 \times 10^{-2} \text{ mm} \end{aligned}$$

26. A capacitor of capacitance 900  $\mu\text{F}$  is charged by a 100 V battery. The capacitor is disconnected from the battery and connected to another uncharged identical capacitor such that one plate of uncharged capacitor connected to positive plate and another plate of uncharged capacitor connected to negative plate of the charged capacitor. The loss of energy in this process is measured as  $x \times 10^{-2}$  J. The value of x is \_\_\_\_\_.

**Answer (225)**

**Sol.**  $U_i = \frac{1}{2} CV^2 = \frac{1}{2} \times 900 \times 10^{-6} \times 100^2 = 4.5$  J

As the other capacitor is identical therefore charge is equally divided and potential difference across the capacitors becomes half. So

$$\begin{aligned} U_f &= \frac{1}{2} 2C \left( \frac{V}{2} \right)^2 = \frac{1}{2} \times 2 \times 900 \times 10^{-6} \left( \frac{100}{2} \right)^2 \\ &= \frac{9}{4} \text{ J} = 2.25 \text{ J} \end{aligned}$$

So, loss in energy  $\Delta U_{\text{loss}} = U_i - U_f$

$$= 2.25 \text{ J}$$

$$= 225 \times 10^{-2} \text{ J}$$

27. A thin uniform of length 2 m, cross sectional area 'A' and density 'd' is rotated about an axis passing through the centre and perpendicular to its length with angular velocity  $\omega$ . If value of  $\omega$  in terms of its rotational kinetic energy E is  $\sqrt{\frac{\alpha E}{Ad}}$  then value of  $\alpha$  is \_\_\_\_\_.

**Answer (3)**

**Sol.** Kinetic energy of rod  $E = \frac{1}{2} \frac{ml^2}{12} \omega^2$

$$\text{or } \omega = \sqrt{\frac{24E}{ml^2}} = \sqrt{\frac{24E}{d \times A \times l^3}}$$


$$\Rightarrow \omega = \sqrt{\frac{24E}{dA l^3}}$$

$$= \sqrt{\frac{3E}{Ad}}$$

So,  $\alpha = 3$




28. In the following circuit, the magnitude of current  $I_1$  is \_\_\_\_\_ A.



**Answer (01.50)**

**Sol.** The indicated diagram shows current flow diagram loops for writing Kirchoff's law are also indicated, writing the equation



$$2I_3 + I_1 + I_3 + I_2 = 5$$

$$\text{or } I_1 + I_2 + 3I_3 = 5 \quad \dots(1)$$

$$I_2 - 5 = 2(I_3 - I_2) + (I_1 + I_3 - I_2)$$

$$\text{or } I_1 - 4I_2 + 3I_3 = -5 \quad \dots(2)$$

$$(I_1 + I_3) + (I_1 + I_3 - I_2) = 2$$

$$\text{or } 2I_1 - I_2 + 2I_3 = 2 \quad \dots(3)$$


$$\text{on solving } I_1 = \frac{3}{2} \text{ A}, I_2 = 2, I_3 = \frac{1}{2} \text{ A}$$

$$= 01.50$$

29. A point source of light is placed at the centre of curvature of a hemispherical surface. The source emits a power of 24 W. The radius of curvature of hemisphere is 10 cm and the inner surface is completely reflecting. The force on the hemisphere due to the light falling on it is \_\_\_\_\_  $\times 10^{-8}$  N.

**Answer (4)**

**Sol.**



$$dA = 2\pi R \sin\theta R d\theta$$

$$= 2\pi R^2 \sin\theta d\theta$$

$$\text{So, } dF = 2 \frac{IdA}{C}$$

$$= \frac{2 \times 24}{4\pi R^2} \times \frac{2\pi R^2 \sin\theta d\theta}{C}$$

$$dF = \frac{24}{C} \sin\theta d\theta$$

This  $dF$  force will be radially outward so the component of this force in vertical direction is

$$dF_v = dF \cos\theta$$

$$\int_0^{F_v} dF_v = \frac{24}{C} \int_0^{\pi/2} \sin\theta \cos\theta d\theta$$

$$= \frac{24}{2C} = \frac{24}{2 \times 3 \times 10^8} = 4 \times 10^{-8} \text{ N}$$

30. The general displacement of a simple harmonic oscillator is  $x = A \sin \omega t$ . Let  $T$  be its time period. The slope of its potential energy ( $U$ ) – time ( $t$ ) curve will

be maximum when  $t = \frac{T}{\beta}$ . The value of  $\beta$  is

**Answer (8)**

$$\text{Sol. } U = \frac{1}{2} m \omega^2 A^2 \sin^2 \omega t$$

$$\text{So, } \frac{dU}{dt} = \frac{m \omega^3 A^2}{2} \sin 2\omega t$$

This value will be maximum when

$$\sin 2\omega t = 1$$

$$\text{or } 2\omega t = \frac{\pi}{2}$$

$$2 \times \frac{2\pi}{T} t = \frac{\pi}{2}$$

$$\Rightarrow t = \frac{T}{8}$$

$$\text{So } \beta = 8$$

**CHEMISTRY****SECTION - A**


**Multiple Choice Questions:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

**Choose the correct answer :**





31. During the qualitative analysis of  $\text{SO}_3^{2-}$  using dilute  $\text{H}_2\text{SO}_4$ ,  $\text{SO}_2$  gas is evolved which turns  $\text{K}_2\text{Cr}_2\text{O}_7$  solution (acidified with dilute  $\text{H}_2\text{SO}_4$ ) :

- (1) red (2) black  
(3) blue (4) green

**Answer (4)**



32. Benzyl isocyanide can be obtained by:


- A.   $\xrightarrow{\text{AgCN}}$
- B.   $\xrightarrow[\text{Aq. KOH}]{\text{CHCl}_3}$
- C.   $\xrightarrow[\text{Aq. KOH}]{\text{CHCl}_3}$
- D.   $\xrightarrow{\text{KCN}}$

Choose the **correct** answer from the options given below :

- (1) A and D (2) Only B  
(3) A and B (4) B and C

**Answer (3)**

**Sol.**



33. In the wet tests for identification of various cations by precipitation, which transition element cation doesn't belong to group IV in qualitative inorganic analysis?

- (1)  $\text{Co}^{2+}$  (2)  $\text{Zn}^{2+}$   
(3)  $\text{Ni}^{2+}$  (4)  $\text{Fe}^{3+}$

**Answer (4)**

**Sol.**  $\text{Fe}^{3+}$  belongs to III<sup>rd</sup> group

34. Amongst the following compounds, which one is an antacid?

- (1) Meprobamate (2) Brompheniramine  
(3) Ranitidine (4) Terfenadine

**Answer (3)**

**Sol.** Ranitidine is not an antacid.

35. The alkaline earth metal sulphate(s) which are readily soluble in water is/are:

- A.  $\text{BeSO}_4$   
B.  $\text{MgSO}_4$   
C.  $\text{CaSO}_4$   
D.  $\text{SrSO}_4$   
E.  $\text{BaSO}_4$

Choose the **correct** answer from the options given below:

- (1) B only (2) A and B  
(3) B and C (4) A only

**Answer (2)**

**Sol.**  $\text{BeSO}_4$  and  $\text{MgSO}_4$  are readily soluble in water.




36. Formation of photochemical smog involves the following reaction in which A, B and C are respectively.

- i.  $\text{NO}_2 \xrightarrow{h\nu} \text{A} + \text{B}$   
ii.  $\text{B} + \text{O}_2 \rightarrow \text{C}$   
iii.  $\text{A} + \text{C} \rightarrow \text{NO}_2 + \text{O}_2$

Choose the correct answer from the options given below:

- (1) O,  $\text{N}_2\text{O}$  and NO (2) NO, O and  $\text{O}_3$   
(3) N,  $\text{O}_2$  and  $\text{O}_3$  (4) O, NO and  $\text{NO}_3^-$


**Answer (2)**



37. Lithium aluminium hydride can be prepared from the reaction of

- (1) LiH and Al(OH)<sub>3</sub>      (2) LiCl and Al<sub>2</sub>H<sub>6</sub>  
 (3) LiCl, Al and H<sub>2</sub>      (4) LiH and Al<sub>2</sub>Cl<sub>6</sub>

**Answer (4)**



38. For OF<sub>2</sub> molecule consider the following:

- A. Number of lone pairs on oxygen is 2.  
 B. FOF angle is less than 104.5°.  
 C. Oxidation state of O is -2.  
 D. Molecule is bent 'V' shaped  
 E. Molecular geometry is linear.

**Correct options are:**

- (1) C, D, E only      (2) B, E, A only  
 (3) A, C, D only      (4) A, B, D only

**Answer (4)**

Sol.








A : No. of lone pairs on oxygen = 2

B : (θ < Bond angle in H<sub>2</sub>O (104.5°))


D : molecule is bent "v" shaped

39. The major products 'A' and 'B', respectively, are



**Answer (4)**


Sol.




40. Match List I with List II

**List I**


**List II**




I. Fittig reaction



II. Wurtz Fittig reaction







III. Finkelstein reaction



Choose the correct answer from the options given below:

- (1) A - II, B - I, C - III, D - IV  
 (2) A - II, B - I, C - IV, D - III  
 (3) A - IV, B - II, C - III, D - I  
 (4) A - III, B - II, C - IV, D - I

**Answer (2)**



41. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A)** : In expensive scientific instruments, silica gel is kept in watch-glasses or in semipermeable membrane bags.

**Reason (R)** : Silica gel adsorbs moisture from air via adsorption, thus protects the instrument from water corrosion (rusting) and / or prevents malfunctioning.

In the light of the above statements, choose the **correct** answer from the options given below :

- (1) Both **(A)** and **(R)** are true but **(R)** is **not** the correct explanation of **(A)**
- (2) **(A)** is false but **(R)** is true
- (3) **(A)** is true but **(R)** is false
- (4) Both **(A)** and **(R)** are true but **(R)** is the correct explanation of **(A)**

**Answer (4)**

Sol. Assertion is correct and Reason is correct explanation of Assertion.

Silica gel adsorbs moisture and thus protects the instrument from water corrosion (rusting) and prevents malfunctioning

42. To inhibit the growth of tumours, identify the compounds used from the following :

- (A) EDTA
- (B) Coordination Compounds of Pt
- (C) D - Penicillamine
- (D) Cis - Platin

Choose the correct answer from the option given below :

- (1) C and D only
- (2) B and D only
- (3) A and B only
- (4) A and C only

**Answer (2)**

Sol. Cis-platin is  $[Pt(NH_3)_2Cl_2]$ ; cis platin and other complexes of pt are used to inhibit the growth of tumours.

43. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A)** : Ketoses give Seliwonoff's test faster than Aldoses.


**Reason (R)** : Ketoses undergo  $\beta$ -elimination followed by formation of furfural.

In the light of the above statements, choose the **correct** answer from the options given below :

- (1) **(A)** is true but **(R)** is false
- (2) Both **(A)** and **(R)** are true and **(R)** is the correct explanation of **(A)**
- (3) **(A)** is false but **(R)** is true
- (4) Both **(A)** and **(R)** are true but **(R)** is **not** the correct explanation of **(A)**

**Answer (1)**

Sol.




This test relies on the principle that, when heated, ketoses are more rapidly dehydrated than Aldoses.

Ketose  $\rightarrow$  Red color formed immediately


Aldose  $\rightarrow$  light pink color formed slowly

44. What is the correct order of acidity of the protons marked A–D in the given compounds?



- (1)  $H_C > H_A > H_D > H_B$     (2)  $H_D > H_C > H_B > H_A$   
 (3)  $H_C > H_D > H_B > H_A$     (4)  $H_C > H_D > H_A > H_B$

**Answer (4)**







$$H_C > H_D > H_A > H_B$$

$H_C$  is hydrogen of carboxylic acid

$H_D$  removal will lead to stable carbanion.

45. Which of the following compounds would give the following set of qualitative analysis?

- (i) Fehling's Test : Positive  
 (ii) Na fusion extract upon treatment with sodium nitroprusside gives a blood red colour but not prussian blue.



**Answer (4)**


**Sol.** Fehling solution is not given by aromatic aldehydes.

1, 2, 3 are aromatic aldehydes

46. In the extraction of copper, its sulphide ore is heated in a reverberatory furnace after mixing with silica to

- (1) Decrease the temperature needed for roasting of  $Cu_2S$   
 (2) Remove calcium as  $CaSiO_3$   
 (3) Separate  $CuO$  as  $CuSiO_3$   
 (4) Remove  $FeO$  as  $FeSiO_3$

**Answer (4)**



47. Match **List I** with **List II**

	List-I (Atomic number)		List-II (Block of periodic table)
A.	37	I.	p-block
B.	78	II.	d-block
C.	52	III.	f-block
D.	65	IV.	s-block

Choose the **correct** answer from the options given below

- (1) A-II, B-IV, C-I, D-III    (2) A-IV, B-III, C-II, D-I  
 (3) A-IV, B-II, C-I, D-III    (4) A-I, B-III, C-IV, D-II

**Answer (3)**

**Sol.**

37 – s-Block

78 – d-Block

52 – p-Block

65 – f-Block

48. Which of the following is correct order of ligand field strength?

- (1)  $NH_3 < en < CO < S^{2-} < C_2O_4^{2-}$   
 (2)  $S^{2-} < NH_3 < en < CO < C_2O_4^{2-}$   
 (3)  $S^{2-} < C_2O_4^{2-} < NH_3 < en < CO$   
 (4)  $CO < en < NH_3 < C_2O_4^{2-} < S^{2-}$

**Answer (3)**

**Sol.** Ligand field strength

$$S^{2-} < C_2O_4^{2-} < NH_3 < en < CO$$

49. Match **List I** with **List II**


	List-I (Molecules/Ions)		List-II (No. of lone pairs of $e^-$ on central atom)
A.	$IF_7$	I.	Three
B.	$ICl_4^-$	II.	One
C.	$XeF_6$	III.	Two
D.	$XeF_2$	IV.	Zero

Choose the **correct** answer from the options given below

- (1) A-II, B-III, C-IV, D-I (2) A-IV, B-I, C-II, D-III  
 (3) A-II, B-I, C-IV, D-III (4) A-IV, B-III, C-II, D-I

**Answer (4)**

- Sol.** (A)  $\text{IF}_7$  – 0 lone pairs  
 (B)  $\text{ICl}_4^-$  – 2 lone pairs  
 (C)  $\text{XeF}_6$  – 1 lone pair  
 (D)  $\text{XeF}_2$  – 3 lone pairs




50. Caprolactam when heated at high temperature in presence of water, gives

- (1) Nylon 6, 6                      (2) Nylon 6  
 (3) Dacron                      (4) Teflon

**Answer (2)**

**Sol.** Caprolactam  $\xrightarrow{\Delta}$  Nylon – 6



**SECTION - B**

**Numerical Value Type Questions:** This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g. 06.25, 07.00, –00.33, –00.30, 30.27, –27.30) using the mouse andw the on-screen virtual numeric keypad in the place designated to enter the answer.

51. A solution containing 2 g of a non-volatile solute in 20 g of water boils at 373.52 K. The molecular mass of the solute is \_\_\_\_\_ g mol<sup>-1</sup>. (Nearest integer)  
 Given, water boils at 373 K,  $K_b$  for water = 0.52 K kg mol<sup>-1</sup>

**Answer (100)**

**Sol.**  $\Delta T_b = K_b \cdot m$   
 $(0.52) = (0.52) (m)$   
 $m = 1 = \frac{2(1000)}{(mw)(20)}$   
 $mw = 100$

52. A 300 mL bottle of soft drink has 0.2 M  $\text{CO}_2$  dissolved in it. Assuming  $\text{CO}_2$  behaves as an ideal gas, the volume of the dissolved  $\text{CO}_2$  at STP is \_\_\_\_\_ mL. (Nearest integer)

Given: At STP, molar volume of an ideal gas is 22.7 L mol<sup>-1</sup>

**Answer (1362)**

**Sol.** Moles = 0.3 × 0.2  
 Volume at STP = 0.3 × 0.2 × 22.7  
 = 1.362 litre  
 = 1362 mL

53. The energy of one mole of photons of radiation of frequency  $2 \times 10^{12}$  Hz in J mol<sup>-1</sup> is \_\_\_\_\_. (Nearest integer)

[Given:  $h = 6.626 \times 10^{-34}$  Js  
 $N_A = 6.022 \times 10^{23}$  mol<sup>-1</sup>]

**Answer (798)**

**Sol.**  $E = nh\nu$   
 $= (6.022 \times 10^{23}) (6.626 \times 10^{-34}) \times (2 \times 10^{12})$   
 $= 798.03$  J  
 $\approx 798$  J

54. Some amount of dichloromethane ( $\text{CH}_2\text{Cl}_2$ ) is added to 671.141 mL of chloroform ( $\text{CHCl}_3$ ) to prepare  $2.6 \times 10^{-3}$  M solution of  $\text{CH}_2\text{Cl}_2$  (DCM). The concentration of DCM is \_\_\_\_\_ ppm (by mass).

Given:  
 Atomic mass : C = 12  
 H = 1  
 Cl = 35.5  
 Density of  $\text{CHCl}_3 = 1.49$  g cm<sup>-3</sup>

**Answer (148)**

**Sol.** Mass of  $\text{CHCl}_3 = 671.141 \times 1.49$   
 $= 1000$  gm  
 $2.6 \times 10^{-3} = \frac{\text{moles of } \text{CH}_2\text{Cl}_2}{0.671141}$

$$\Rightarrow \text{moles of CH}_2\text{Cl}_2 = 1.74496 \times 10^{-3}$$


$$\text{mass of CH}_2\text{Cl}_2 = 148.32 \times 10^{-3} \text{ gm}$$

$$\text{Composition of CH}_2\text{Cl}_2 = \frac{148.32 \times 10^{-3}}{1000} \times 10^6$$

$$= 148.32 \text{ ppm}$$

$$\approx 148$$

55. Consider the cell



When the potential of the cell is 0.712 V at 298 K, the ratio  $[\text{Fe}^{2+}] / [\text{Fe}^{3+}]$  is \_\_\_\_\_.

(Nearest integer)


Given:  $\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$ ,  $E^\ominus_{\text{Fe}^{3+}, \text{Fe}^{2+}} | \text{Pt} = 0.771$

$$\frac{2.303 RT}{F} = 0.06 \text{ V}$$

**Answer (10)**

**Sol.** Anode  $\text{H}_2 \rightarrow 2\text{H}^+ + 2\text{e}^-$

Cathode  $(\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+}) \times 2$



$$E_{\text{cell}} = E_{\text{cell}}^\ominus - \frac{0.059}{2} \log \left( \frac{[\text{Fe}^{2+}]^2}{[\text{Fe}^{3+}]^2} \right)$$

$$0.712 = 0.771 - 0.059 \log \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]}$$

$$-0.059 = -0.059 \log \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]}$$

$$\frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]} = 10$$

56. If compound A reacts with B following first order kinetics with rate constant  $2.011 \times 10^{-3} \text{ s}^{-1}$ . The time taken by A (in seconds) to reduce from 7 g to 2 g will be \_\_\_\_\_. (Nearest Integer)

[ $\log 5 = 0.698$ ,  $\log 7 = 0.845$ ,  $\log 2 = 0.301$ ]

**Answer (623)**

$$\text{Sol. } t = \frac{2.303}{k} \log \frac{C_0}{C_t}$$

$$= \frac{2.303}{2.011 \times 10^{-3}} \log \frac{7}{2}$$

$$= \frac{2.303 \times 10^3}{2.011} (.845 - .301)$$

$$= 622.99$$

$$\approx 623 \text{ sec.}$$


57. When 2 litre of ideal gas expands isothermally into vacuum to a total volume of 6 litre, the change in internal energy is \_\_\_\_\_ J. (Nearest integer)

**Answer (0)**

**Sol.** For isothermal process of an ideal gas;  $\Delta E = 0$

58. The number of electrons involved in the reduction of permanganate of manganese dioxide in acidic medium is \_\_\_\_\_.

**Answer (3)**



59. 600 mL of 0.01 M HCl is mixed with 400 mL of 0.01 M  $\text{H}_2\text{SO}_4$ . The pH of the mixture is \_\_\_\_\_  $\times 10^{-2}$ . (Nearest integer)

[Given  $\log 2 = 0.30$

$\log 3 = 0.48$

$\log 5 = 0.69$

$\log 7 = 0.84$

$\log 11 = 1.04$ ]

**Answer (186)**

$$\text{Sol. } [\text{H}^+] = \frac{6+8}{1000} = 14 \times 10^{-3}$$

$$\text{pH} = 3 - \log 14$$

$$= 3 - .3 - .84$$

$$= 1.86 = 186 \times 10^{-2}$$

60. A trisubstituted compound 'A',  $\text{C}_{10}\text{H}_{12}\text{O}_2$  gives neutral  $\text{FeCl}_3$  test positive. Treatment of compound 'A' with NaOH and  $\text{CH}_3\text{Br}$  gives  $\text{C}_{11}\text{H}_{14}\text{O}_2$ , with hydroiodic acid gives methyl iodide and with hot conc. NaOH gives a compound B,  $\text{C}_{10}\text{H}_{12}\text{O}_2$ . Compound 'A' also decolorises alkaline  $\text{KMnO}_4$ . The number of  $\pi$  bond/s present in the compound 'A' is \_\_\_\_\_.

**Answer (4)**

**Sol.** A :  $\text{C}_{10}\text{H}_{12}\text{O}_2$

$$\text{DU of A} = \frac{22-12}{2} = 5$$

1 DU is due to Ring (Benzene ring)

4  $\pi$ -bonds will be there

(3  $\pi$ -bonds in ring and 1  $\pi$ -bond outside ring) as it decolorises alkaline  $\text{KMnO}_4$ .

**MATHEMATICS****SECTION - A**

**Multiple Choice Questions:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

**Choose the correct answer :**

61. If  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  are three non-zero vectors and  $\hat{n}$  is a unit vector perpendicular to  $\vec{c}$  such that  $\vec{a} = \alpha\vec{b} - \hat{n}$ , ( $\alpha \neq 0$ ) and  $\vec{b} \cdot \vec{c} = 12$ , then  $|\vec{c} \times (\vec{a} \times \vec{b})|$  is equal to

- (1) 9 (2) 6  
(3) 12 (4) 15

**Answer (3)**

**Sol.**  $\hat{n} = \alpha\vec{b} - \vec{a}$

$$\begin{aligned} \vec{c} \times (\vec{a} \times \vec{b}) &= (\vec{c} \cdot \vec{b})\vec{a} - (\vec{c} \cdot \vec{a})\vec{b} \\ &= 12\vec{a} - (\vec{c} \cdot (\alpha\vec{b} - \hat{n}))\vec{b} \\ &= 12\vec{a} - (12\alpha - 0)\vec{b} \\ &= 12(\vec{a} - \alpha\vec{b}) \end{aligned}$$

$$\therefore |\vec{c} \times (\vec{a} \times \vec{b})| = 12$$

62. If the coefficient of  $x^{15}$  in the expansion of  $\left(ax^3 + \frac{1}{bx^{1/3}}\right)^{15}$  is equal to the coefficient of  $x^{-15}$  in

the expansion of  $\left(ax^{1/3} - \frac{1}{bx^3}\right)^{15}$ , where  $a$  and  $b$

are positive real numbers, then for each such ordered pair  $(a, b)$

- (1)  $ab = 1$  (2)  $a = b$   
(3)  $a = 3b$  (4)  $ab = 3$

**Answer (1)**

**Sol.** For  $\left(ax^3 + \frac{1}{bx^{1/3}}\right)^{15}$

$$T_{r+1} = {}^{15}C_r (ax^3)^{15-r} \left(\frac{1}{bx^{1/3}}\right)^r$$

$$\therefore \text{For } x^{15} \rightarrow 3(15-r) - \frac{r}{3} = 15$$

$$\Rightarrow 30 = \frac{10r}{3} \Rightarrow r = 9$$

Similarly, for  $\left(ax^3 - \frac{1}{bx^3}\right)^{15}$

$$T_{r+1} = {}^{15}C_r \left(ax^3\right)^{15-r} \left(\frac{1}{bx^3}\right)^r$$

$$\therefore \text{For } x^{-15} \rightarrow \frac{15-r}{3} - 3r = -15 \Rightarrow r = 6$$

$$\therefore {}^{15}C_9 \frac{a^6}{b^9} = {}^{15}C_6 \frac{a^9}{b^6} \Rightarrow ab = 1$$

63. The number of points on the curve  $y = 54x^5 - 135x^4 - 70x^3 + 180x^2 + 210x$  at which the normal lines are parallel to  $x + 90y + 2 = 0$  is

- (1) 4 (2) 3  
(3) 2 (4) 0

**Answer (1)**

**Sol.**  $y' = 270x^4 - 540x^3 - 210x^2 + 360x + 210$

$$\text{Slope of normal} = -\frac{1}{90}$$

$$\therefore \text{Slope of tangent} = 90$$

$\therefore$  Number of normal will be number of solutions of

$$270x^4 - 540x^3 - 210x^2 + 360x + 210 = 90$$

$$\Rightarrow 9x^4 - 18x^3 - 7x^2 + 12x + 4 = 0$$

$$\therefore x = 1, 2, -\frac{1}{3}, -\frac{2}{3} \text{ are roots}$$

64. The line  $l_1$  passes through the point  $(2, 6, 2)$  and is perpendicular to the plane  $2x + y - 2z = 10$ . Then the shortest distance between the line  $l_1$  and the line

$$\frac{x}{2} - \frac{y+4}{-3} = \frac{z}{2} \text{ is}$$

- (1)  $\frac{19}{3}$  (2)  $\frac{13}{3}$   
(3) 9 (4) 7

**Answer (3)**



Sol. Equation of  $l_1 = \frac{x-2}{2} = \frac{y-6}{1} = \frac{z-2}{-2}$

Shortest distance with  $\frac{x+1}{2} = \frac{y+4}{-3} = \frac{z}{2}$  is

$$\text{S.d} = \frac{\begin{vmatrix} 3 & 10 & 2 \\ 2 & 1 & -2 \\ 2 & -3 & 2 \end{vmatrix}}{\begin{vmatrix} -4\hat{i} & -8\hat{j} & -8\hat{k} \end{vmatrix}} = \frac{|(-12) - 10(8) + 2(-8)|}{12}$$


= 9 units

65. Let  $y = x + 2$ ,  $4y = 3x + 6$  and  $3y = 4x + 1$  be three tangent lines to the circle  $(x - h)^2 + (y - k)^2 = r^2$ . Then  $h + k$  is equal to

- (1)  $5(1 + \sqrt{2})$                       (2) 5  
 (3) 6                                      (4)  $5\sqrt{2}$

Answer (2)

Sol.



$$(h, k) = \left( \frac{5 \cdot 5 + 5(-2) + 14\sqrt{2}}{10 + 7\sqrt{2}}, \frac{35 + 21\sqrt{2}}{10 + 7\sqrt{2}} \right)$$

$$h + k = \frac{50 + 35\sqrt{2}}{10 + 7\sqrt{2}} = 5$$

66. Among the statements :

- (S1)  $((p \vee q) \Rightarrow r) \Leftrightarrow (p \Rightarrow r)$   
 (S2)  $((p \vee q) \Rightarrow r) \Leftrightarrow ((p \Rightarrow r) \vee (q \Rightarrow r))$   
 (1) only (S1) is a tautology  
 (2) only (S2) is a tautology  
 (3) neither (S1) nor (S2) is a tautology  
 (4) both (S1) and (S2) are tautologies

Answer (3)

Sol.  $S_1 : ((p \vee q) \Rightarrow r) \Leftrightarrow (p \Rightarrow r)$

$$S_2 : ((p \vee q) \Rightarrow r) \Leftrightarrow ((p \Rightarrow r) \vee (q \Rightarrow r))$$

In  $S_1$  : If  $p = F$ ,  $q = T$ ,  $r = F$  then  $S_1$  is false

In  $S_2$  : if  $P = T$ ,  $q = F$ ,  $r = F$  then  $S_2$  is false

$\therefore$  Neither  $S_1$  nor  $S_2$  is a tautology

67. The coefficient of  $x^{301}$  in  $(1+x)^{500} + x(1+x)^{499} + x^2(1+x)^{498} + \dots + x^{500}$  is

- (1)  $^{501}C_{302}$                       (2)  $^{501}C_{200}$   
 (3)  $^{500}C_{300}$                       (4)  $^{500}C_{301}$

Answer (2)

Sol.  $^{500}C_{301} + ^{499}C_{300} + ^{498}C_{299} + \dots + ^{199}C_0$   
 $= ^{500}C_{199} + ^{499}C_{199} + ^{498}C_{199} + \dots + ^{199}C_{89}$   
 $= ^{501}C_{200}$

68. The minimum number of elements that must be added to the relation  $R = \{a, b\}, \{b, c\}$  on the set  $\{a, b, c\}$  so that it becomes symmetric and transitive is

- (1) 7                                      (2) 3  
 (3) 5                                      (4) 4

Answer (1)

Sol. For symmetric  $(b, a), (c, b) \in R$

For transitive  $(a, c) \in R$

$$\Rightarrow (c, a) \in R$$

$$\therefore (a, b), (b, a) \in R$$

$$\Rightarrow (a, a) \in R$$

$$(b, c), (c, b) \in R$$

$$\Rightarrow (b, b) \in R, (c, c) \in R$$

7 elements must be added

69. If  $P(h, k)$  be a point on the parabola  $x = 4y^2$ , which is nearest to the point  $Q(0, 33)$ , then the distance of  $P$  from the directrix of the parabola  $y^2 = 4(x + y)$  is equal to

- (1) 4                                      (2) 6  
 (3) 8                                      (4) 2

Answer (2)

Sol. Equation of normal

$$y = -tx + 2 \cdot \frac{1}{16}t + \frac{1}{16}t^3$$

$$33 = \frac{t}{8} + \frac{t^3}{16}$$

$$\Rightarrow t^3 + 2t = 528$$

$$t = 8$$

$$(at^2, 2at) = (4, 1)$$

Distance from  $x = -2$

70. If  $[t]$  denotes the greatest integer  $\leq t$ , then the value

$$\text{of } \frac{3(e-1)}{e} \int_1^2 x^2 e^{x^3} dx \text{ is}$$

- (1)  $e^7 - 1$                       (2)  $e^8 - 1$   
 (3)  $e^9 - e$                       (4)  $e^8 - e$

**Answer (4)**

$$\begin{aligned} \text{Sol. } I &= \frac{3(e-1)}{e} \int_1^2 x^2 e^{[x]+[x^3]} dx \\ &= \frac{3(e-1)}{e} \int_1^2 x^2 e^{1+[x^3]} dx \quad (\because [x]=1 \text{ when } x \in (1,2)) \end{aligned}$$

$$3(e-1) \int_1^2 x^2 e^{x^3} dx$$

Let  $x^3 = t$

$$I = (e-1) \int_1^8 e^{[t]} dt$$

$$= (e-1)(e^1 + e^2 + e^3 + \dots + e^7)$$

$$(e-1)e \frac{(e^7 - 1)}{e - 1}$$

$$e^8 - e$$

71. Let a unit vector  $\widehat{OP}$  makes angles  $\alpha, \beta, \gamma$  with the positive directions of the co-ordinate axes  $OX, OY, OZ$  respectively, where  $\beta \in \left(0, \frac{\pi}{2}\right)$ . If  $\widehat{OP}$  is perpendicular to the plane through points  $(1, 2, 3), (2, 3, 4)$  and  $(1, 5, 7)$ , then which one of the following is true?

- (1)  $\alpha \in \left(0, \frac{\pi}{2}\right)$  and  $\gamma \in \left(0, \frac{\pi}{2}\right)$   
 (2)  $\alpha \in \left(\frac{\pi}{2}, \pi\right)$  and  $\gamma \in \left(0, \frac{\pi}{2}\right)$   
 (3)  $\alpha \in \left(\frac{\pi}{2}, \pi\right)$  and  $\gamma \in \left(\frac{\pi}{2}, \pi\right)$   
 (4)  $\alpha \in \left(0, \frac{\pi}{2}\right)$  and  $\gamma \in \left(\frac{\pi}{2}, \pi\right)$

**Answer (3)**

**Sol.** Let  $A \equiv (1, 2, 3), B \equiv (2, 3, 4), C \equiv (1, 5, 7)$

$$\vec{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} i & j & k \\ 1 & 1 & 1 \\ 0 & 3 & 4 \end{vmatrix}$$

$$= \hat{i} - 4\hat{j} + 3\hat{k}$$

$$\widehat{OP} = \frac{\pm(\hat{i} - 4\hat{j} + 3\hat{k})}{\sqrt{26}}$$

Since  $\cos\beta > 0$ , take  $-$  sign

$$\widehat{OP} = \frac{-\hat{i} - 4\hat{j} + 3\hat{k}}{\sqrt{26}}$$

$$\Rightarrow \cos\alpha < 0, \cos\gamma < 0$$

$$\alpha, \gamma \in \left(\frac{\pi}{2}, \pi\right)$$

72. Suppose  $f: \mathbb{R} \rightarrow (0, \infty)$  be a differentiable function such that  $5f(x+y) = f(x) \cdot f(y), \forall x, y \in \mathbb{R}$ . If  $f(3) = 320$ ,

then  $\sum_{n=0}^5 f(n)$  is equal to

- (1) 6825                              (2) 6525  
 (3) 6875                              (4) 6575

**Answer (1)**

**Sol.**  $5f(x+y) = f(x) \cdot f(y)$

$$5f(3) = f(1) \cdot f(2)$$

$$5f(2) = (f(1))^2 \quad f(10) = 5$$

$$f(1) = 20$$

$$f(1) \cdot \frac{(f(1))^2}{5} = 1600$$

$$\begin{aligned} \sum_{n=0}^5 f(n) &= f(0) + 20 + 80 + 320 + 1280 + 5120 \\ &= 1750 + 5120 \\ &= 6875 \end{aligned}$$

73. Let  $A = \begin{pmatrix} m & n \\ p & q \end{pmatrix}$ ,  $d = |A| \neq 0$  and  $|A - d(\text{Adj } A)| = 0$ .

Then

$$(1) (1+d)^2 = m^2 + q^2 \quad (2) (1+d)^2 = (m+q)^2$$

$$(3) 1+d^2 = (m+q)^2 \quad (4) 1+d^2 = m^2 + q^2$$

**Answer (2)**

Sol.  $\begin{vmatrix} -d & \begin{vmatrix} q & -n \\ -p & m \end{vmatrix} \end{vmatrix} =$

$$\begin{vmatrix} m - qd & n(1 + d) \\ p(1 + d) & q - md \end{vmatrix} = 0$$

$$(m - qd)(q - md) = np(1 + d)^2$$

$$mq - (q^2 + m^2)d + qmd^2 = np(1 + d^2) + 2npd$$

$$d^2(mq - np) + 1(mq - np) = (2np + m^2 + q^2)d$$

$$(d^2 + 1)(mq - np) = (2np + m^2 + q^2)d$$

$$d^2 + 1 = 2np + m^2 + q^2$$

$$2d = 2mq - 2np$$

$$\Rightarrow (1 + d)^2 = (m + q)^2$$

74. If  $\tan 15^\circ + \frac{1}{\tan 75^\circ} + \frac{1}{\tan 105^\circ} + \tan 195^\circ = 2a$ , then

the value of  $\left(a + \frac{1}{a}\right)$  is

(1) 4 (2) 2

(3)  $4 - 2\sqrt{3}$  (4)  $5 - \frac{3}{2}\sqrt{3}$

**Answer (1)**

Sol.  $\tan 15^\circ + \tan 15^\circ - \tan 15^\circ + \tan 15^\circ = 2\tan 15^\circ$

$$= 2(2 - \sqrt{3}) = 2a \Rightarrow a = 2 - \sqrt{3}$$

$$\therefore \frac{1}{a} + a \Rightarrow (2 + \sqrt{3}) + (2 - \sqrt{3}) = 4$$

75. Let the system of linear equations

$$x + y + kz = 2$$

$$2x + 3y - z = 1$$

$$3x + 4y + 2z = k$$

have infinitely many solutions. Then the system

$$(k + 1)x + 2(k - 1)y = 7$$

$$(2k + 1)x + (k + 5)y = 10$$

has :

(1) Unique solution satisfying  $x + y = 1$

(2) Unique solution satisfying  $x - y = 1$

(3) Infinitely many solutions

(4) No solution

**Answer (1)**

Sol.  $x + y + kz = 2 \dots(1)$

$2x + 3y - z = 1 \dots(2)$

$3x + 4y + 2z = k \dots(3)$

(1) + (2)

$$3x + 4y + z(k - 1) = 3$$

Comparing with (3)

$$k = 3$$

$$\text{Now, } 4x + 5y = 7 \Rightarrow 3x + 3y = 3$$

$$7x + 8y = 10$$

$$\text{as } \frac{4}{7} \neq \frac{5}{8}$$

$\therefore$  unique solution satisfying  $x + y = 1$

76. If the solution of the equation  $\log_{\cos x} \cot x + 4 \log_{\sin x}$

$$\tan x = 1, x \in \left(0, \frac{\pi}{2}\right), \text{ is } \sin^{-1}\left(\frac{\alpha + \sqrt{\beta}}{2}\right), \text{ where } \alpha, \beta$$

are integers, then  $\alpha + \beta$  is equal to

(1) 6

(2) 5

(3) 4

(4) 3

**Answer (3)**

Sol.  $\log_{\cos x} \cot x + 4 \log_{\sin x} \tan x = 1$

$$\Rightarrow \log_{\cos x} \cot x - 4 \log_{\sin x} \cot x = 1$$

$$\Rightarrow 1 - \log_{\cos x} \sin x - 4 - 4 \log_{\sin x} \cos x = 1$$

Let  $\log_{\cos x} \sin x = t$

$$t + \frac{4}{t} = 4$$

$$\Rightarrow t = 2$$

$$\sin x = \cos^2 x$$

$$\Rightarrow \sin x = 1 - \sin^2 x$$

$$\Rightarrow \sin^2 x + \sin x - 1 = 0$$

$$\Rightarrow \sin x = \frac{-1 \pm \sqrt{5}}{2}$$

as  $x \in \left(0, \frac{\pi}{2}\right)$

$$\sin x = \frac{\sqrt{5} - 1}{2}$$

$$x = \sin^{-1}\left(\frac{-1 + \sqrt{5}}{2}\right)$$

$$\Rightarrow \alpha = -1, \beta = 5$$


$$\alpha + \beta = 4$$

77. A straight line cuts off the intercepts  $OA = a$  and  $OB = b$  on the positive direction of x-axis and y-axis respectively. If the perpendicular from origin  $O$  to this line makes an angle of  $\frac{\pi}{6}$  with positive direction of y-axis and the area of  $\triangle OAB$  is  $\frac{98}{3}\sqrt{3}$ , then  $a^2 - b^2$  is equal to

- (1) 196 (2)  $\frac{196}{3}$   
 (3)  $\frac{392}{3}$  (4) 98

**Answer (3)**

Sol.  $\frac{1}{2}ab = \frac{98\sqrt{3}}{3}$



$\Rightarrow \sqrt{3}ab = 196 \dots(i)$   
 $OP = OB \cos 30^\circ = OA \cos 60^\circ$   
 $\Rightarrow \frac{b\sqrt{3}}{2} = \frac{a}{2}$   
 $\Rightarrow \sqrt{3}b = a \dots(ii)$

By (i) and (ii)  
 $a^2 = 196$   
 $a = 14$

$b^2 = \frac{a^2}{3}$

$a^2 - b^2 = \frac{2a^2}{3} = \frac{392}{3}$

78. If  $a_n = \frac{-2}{4n^2 - 16 + 5}$ , then  $a_1 + a_2 + \dots + a_{25}$  is equal to

- (1)  $\frac{49}{138}$  (2)  $\frac{52}{147}$   
 (3)  $\frac{51}{144}$  (4)  $\frac{50}{141}$

**Answer (4)**

Sol.  $\sum_{i=1}^{25} a_i = \sum \frac{2}{4^2 - 16n + 15} = \sum \frac{-2}{(2n-5)(2n-3)}$   
 $= \sum_{i=1}^{25} \left( \frac{1}{2n-3} - \frac{1}{2n-5} \right)$   
 $= \left[ \left( \frac{1}{-1} - \frac{1}{-3} \right) + \left( \frac{1}{1} - \frac{1}{-1} \right) + \left( \frac{1}{3} - \frac{1}{1} \right) \dots \right]$   
 $= \frac{1}{2(25)-3} + \frac{1}{3} = \frac{50}{141}$

79. Let the solution curve  $y = y(x)$  of the differential equation

$\frac{dy}{dx} - \frac{3x^5 \tan^{-1}(x^3)}{(1+x^6)^{3/2}} y = 2x \exp \left\{ \frac{x^3 - \tan^{-1} x^3}{\sqrt{1+x^6}} \right\}$

pass through the origin. Then  $y(1)$  is equal to

- (1)  $\exp\left(\frac{1-\pi}{4\sqrt{2}}\right)$  (2)  $\exp\left(\frac{4-\pi}{4\sqrt{2}}\right)$   
 (3)  $\exp\left(\frac{4+\pi}{4\sqrt{2}}\right)$  (4)  $\exp\left(\frac{\pi-4}{4\sqrt{2}}\right)$

**Answer (2)**

Sol.  $\frac{dy}{dx} - \frac{3x^5 \tan^{-1}(x^3)}{(1+x^6)^{3/2}} y = 2x \exp \left\{ \frac{x^3 - \tan^{-1} x^3}{\sqrt{1+x^6}} \right\}$   
 $- \int \frac{3x^5 \tan^{-1}(x^3)}{(1+x^6)^{3/2}} dx$   
 $IF = e^{\dots}$

Let  $\tan^{-1} x^3 = t \Rightarrow \frac{3x^2}{1+x^6} dx = dt$

$\Rightarrow IF = e^{-\int \frac{\tan t}{\sec t} t dt} = e^{-\int \sin t \cdot t dt} = e^{t \cos t - \sin t}$

$\Rightarrow IF = e^{\frac{\tan^{-1}(x^3)}{\sqrt{1+x^6}} - \frac{x^3}{\sqrt{1+x^6}}}$

$\therefore$  Solution is

$y \cdot e^{\frac{\tan^{-1} x^3}{\sqrt{1+x^6}} - \frac{x^3}{\sqrt{1+x^6}}} = \int 2x dx + c$

$\Rightarrow y \cdot e^{\frac{\tan^{-1} x^3 - x^3}{\sqrt{1+x^6}}} = x^2 + c$

$y(0) = 0 \Rightarrow c = 0$

$$x = 1$$

$$y \cdot e^{\frac{\pi-1}{4\sqrt{2}}} = 1$$

$$\Rightarrow y = e^{\frac{1-\pi}{4\sqrt{2}}}$$

$$\Rightarrow y = e^{\frac{4-\pi}{4\sqrt{2}}}$$

80. If an unbiased die, marked with  $-2, -1, 0, 1, 2, 3$  on its faces, is thrown five times, then the probability that product of the outcomes is positive is:

(1)  $\frac{881}{2592}$

(2)  $\frac{440}{2592}$

(3)  $\frac{27}{288}$

(4)  $\frac{521}{2592}$

**Answer (4)**

**Sol.**  ${}^5C_0 \times 3^5 = 243$

$${}^5C_2 \times 2^2 \times 3^3 = 1080$$

$${}^5C_4 \times 2^4 \cdot 3 = 240$$

$\therefore$  required probability

$$= \frac{243 + 1080 + 240}{6 \times 6 \times 6 \times 6 \times 6} = \frac{521}{2592}$$

**SECTION - B**

**Numerical Value Type Questions:** This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g. 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

81. Let  $S = \{1, 2, 3, 4, 5, 6\}$ . Then the number of one-one functions  $f: S \rightarrow P(S)$ , where  $P(S)$  denote the power set of  $S$ , such that  $f(n) \subset f(m)$  where  $n < m$  is \_\_\_\_\_.


**Answer (3240)**

**Sol.**  $\therefore S = \{1, 2, 3, 4, 5, 6\}$

and  $P(S) = \{\phi, \{1\}, \{2\}, \dots, \{1, 2, 3, 4, 5, 6\}\}$

$f(n)$  corresponding a set having  $m$  elements which belongs to  $P(S)$ , should be a subset of  $f(n+1)$ , so  $f(n+1)$  should be a subset of  $P(S)$  having at least  $m+1$  elements.

Now if  $f(1)$  has one element then  $f(2)$  has 3,  $f(3)$  has 3 and so on and  $f(6)$  has 6 elements. Total number of possible functions =  $6! = 720 \dots (1)$  if  $f(1)$  has no elements (i.e. null set  $\phi$ ) then



Each index number represents the number of elements in respective rows

Taking every series of arrow and counting number of such possible functions (sets)

$$= {}^6C_2 \times {}^4C_1 \times {}^3C_1 \times {}^2C_1 + {}^6C_1 \times {}^5C_2 \times {}^3C_1 \times {}^2C_1 + {}^6C_1 \times {}^5C_1 \times {}^4C_2 \times {}^2C_1 + {}^6C_1 \times {}^5C_1 \times {}^4C_1 \times {}^3C_2 + {}^6C_1 \times {}^5C_1 \times {}^4C_1 \times {}^3C_1 \times {}^2C_1 = 2520 \dots (2)$$

From (1) and (2) : Total number of functions =  $2520 + 720 = 3240$

82. Number of 4-digit numbers (the repetition of digits is allowed) which are made using the digits 1, 2, 3 and 5, and are divisible by 15, is equal to \_\_\_\_\_

**Answer (21)**

**Sol.** Digits 1, 2, 3, 5 and number should be divisible by 15 (i.e., divisible by both 3 and 5)

So,

$$\dots (5)$$

**Case-I:**  $5 \rightarrow 1, 2 \rightarrow 1, 1 \rightarrow 2 = \frac{3!}{2!} = 3$

$$5 \rightarrow 1, 3 \rightarrow 1, 2 \rightarrow 2 = \frac{3!}{2!} = 3$$

$$5 \rightarrow 1, 3 \rightarrow 2, 1 \rightarrow 1 = \frac{3!}{2!} = 3$$

**Case-II:**  $5 \rightarrow 2, 3 \rightarrow 1, 2 \rightarrow 1 = 3! = 6$

$$5 \rightarrow 2, 1 \rightarrow 2 = \frac{3!}{2!} = 3$$

**Case-III:**  $5 \rightarrow 3, 3 \rightarrow 1 = \frac{3!}{2!} = 3$

$\therefore$  Total no. = 21

83. If  $\lambda_1 < \lambda_2$  are two values of  $\lambda$  such that the angle between the planes  $P_1 : \vec{r} \cdot (3\hat{i} - 5\hat{j} + \hat{k}) = 7$  and

$P_2 : \vec{r} \cdot (\lambda\hat{i} + \hat{j} - 3\hat{k}) = 9$  is  $\sin^{-1}\left(\frac{2\sqrt{6}}{5}\right)$ , then the

square of the length of perpendicular from the point  $(38\lambda_1, 10\lambda_2, 2)$  to the plane  $P_1$  is \_\_\_\_\_.

**Answer (315)**

**Sol.**  $P_1 : \vec{r} \cdot (3\hat{i} - 5\hat{j} + \hat{k}) = 7$

$P_2 : \vec{r} \cdot (\lambda\hat{i} + \hat{j} - 3\hat{k}) = 9$

Let angle between  $P_1$  and  $P_2$  is  $\theta$

Then  $\cos\theta = \frac{3\lambda - 5 - 3}{\sqrt{35}\sqrt{\lambda^2 + 1}}$

But  $\sin\theta = \frac{2\sqrt{6}}{5}$

$\therefore \frac{(3\lambda - 8)^2}{35(\lambda^2 + 1)} = 1 - \frac{24}{25}$

$\Rightarrow 5(9\lambda^2 + 64 - 48\lambda) = 7\lambda^2 + 70$

$\Rightarrow 38\lambda^2 - 240\lambda + 250 = 0$

$\Rightarrow 19\lambda^2 - 120\lambda + 125 = 0$

$\Rightarrow (19\lambda - 25)(\lambda - 5) = 0$

$\therefore \lambda = \frac{25}{19}, \lambda_2 = 5$

So, point  $(50, 50, 2)$

$\therefore d = \frac{|150 - 250 + 2 - 7|}{\sqrt{35}} = 315$

84. Let  $\sum_{n=0}^{\infty} \frac{n^3((2n)!) + (2n-1)(n!)}{(n!)((2n)!)} = ae + \frac{b}{e} + c$ ,

where  $a, b, c \in \mathbb{Z}$  and  $e = \sum_{n=0}^{\infty} \frac{1}{n!}$ . Then  $a^2 - b + c$  is

equal to \_\_\_\_\_.

**Answer (26)**

**Sol.**  $\sum_{n=0}^{\infty} \frac{n^3(2n!) + (2n-1)(n!)}{n! \cdot (2n)!}$

$\sum_{n=0}^{\infty} \frac{3}{n!} + \frac{2n-1}{2n!}$

$= \sum_{n=0}^{\infty} \frac{3}{(n-2)!} + \frac{1}{(n-3)!} + \frac{1}{(n-1)!} + \frac{1}{(2n-1)!} - \frac{1}{(2n)!}$

$= 3e + e + e - \frac{1}{e}$

$= 5e - \frac{1}{e}$

$\therefore a = 5, b = -1, c = 0$

$\therefore a^2 - b + c = 26$

85. Let  $z = 1 + i$  and  $z_1 = \frac{1 + \bar{z}}{\bar{z}(1-z) + \frac{1}{z}}$ . Then  $\frac{12}{\pi} \arg(z_1)$

is equal to \_\_\_\_\_.

**Answer (09)**

**Sol.**  $z = 1 + i$

$z_1 = \frac{1 + \bar{z}}{\bar{z}(1-z) + \frac{1}{z}}$

$= \frac{z(1 + \bar{z})}{|z|^2(1-z) + 1}$

$= \frac{(1+i)(1+i(1-i))}{2(1-1-i) + 1}$

$z_1 = 1 - i$

$\arg z_1 = \tan^{-1}\left(\frac{-1}{1}\right) = \frac{3\pi}{4}$

$\frac{12}{\pi} \arg(z_1) = \frac{3\pi}{4} \cdot \frac{12}{\pi} = 9$

86. Let  $f^1(x) = \frac{3x+2}{2x+3}, x \in R - \left\{-\frac{3}{2}\right\}$

For  $n \geq 2$ , define  $f^n(x) = f^1$  of  $f^{n-1}(x)$ .

If  $f^5(x) = \frac{ax+b}{bx+a}$ ,  $\gcd(a, b) = 1$ , then  $a + b$  is equal to \_\_\_\_\_.

**Answer (3125)**

**Sol.**  $f^1(x) = \frac{3x+2}{2x+3}, x \in R - \left\{-\frac{3}{2}\right\}$

$f^5(x) = f_o f_o f_o f_o f(x)$

$f_o f(x) = \frac{13x+12}{12x+13}$

$f_o f_o f_o f_o f(x) = \frac{1563x+1562}{1562x+1563}$

$\equiv \frac{ax+b}{bx+a}$

$\therefore a = 1563, b = 1562 = 3125$

87.  $\lim_{x \rightarrow 0} \frac{48}{x^4} \int_0^x \frac{t^3}{t^6 + 1} dt$  is equal to \_\_\_\_\_.

**Answer (12)**

**Sol.**  $\lim_{x \rightarrow 0} \frac{48}{x^4} \int_0^x \frac{t^3}{t^6 + 1} dt$

$$\lim_{x \rightarrow 0} \frac{48}{4x^3} \cdot \left( \frac{x^3}{x^6 + 1} \right)$$

$$\lim_{x \rightarrow 0} \frac{12}{x^6 + 1} = 12$$

88. The mean and variance of 7 observations are 8 and 16 respectively. If one observation 14 is omitted and  $a$  and  $b$  are respectively mean and variance of remaining 6 observation, then  $a + 3b - 5$  is equal to \_\_\_\_\_.

**Answer (37)**

**Sol.**  $\sum x_i = 7 \times 8 = 56$

$$\frac{x^2}{n} - \left( \frac{\sum x_i}{n} \right)^2 = 16$$

$$\frac{\sum x_i^2}{7} - 64 = 16$$

$$\sum x_i^2 = 560$$

When 14 is omitted

$$\sum x_i = 56 - 14 = 42$$

$$\text{New mean} = a = \frac{\sum x_i}{6} = 7$$

$$\sum x_i^2 = 560 - 196 = 364$$

$$\text{new variance, } b = \frac{\sum x_i^2}{6} - \left( \frac{\sum x_i}{6} \right)^2$$

$$= \frac{364}{6} - 49 = \frac{35}{3}$$

$$3b = 35$$

$$a + 3b - 5 = 7 + 35 - 5 = 37$$

89. If the equation of the plane passing through the point  $(1, 1, 2)$  and perpendicular to the line  $x - 3y + 2z - 1 = 0 = 4x - y + z$  is  $Ax + By + Cz = 1$ , then  $140(C - B + A)$  is equal to \_\_\_\_\_.

**Answer (15)**

**Sol.** Line of intersection of the planes  $x - 3y + 2z - 1 = 0$  and  $4x - y + z = 0$  is normal  $(\vec{n})$  to the required plane.

$$\vec{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 2 \\ 4 & -1 & 1 \end{vmatrix} = -\hat{i} + 7\hat{j} + 11\hat{k}$$

Equation of plane is

$$-x + 7y + 11z = \lambda$$

It passes through  $(1, 1, 2)$

$$\therefore \lambda = 28$$

So, the plane is

$$-x + 7y + 11z = 28$$

$$\Rightarrow \frac{-}{28}x + \frac{7}{28}y + \frac{11}{28}z = 1$$


$$A = \frac{-}{28}, B = \frac{7}{28}, C = \frac{11}{28}$$

$$140(C - B + A) = 15$$

90. Let  $\alpha$  be the area of the larger region bounded by the curve  $y^2 = 8x$  and the line  $y = x$  and  $x = 2$ , which lies in the first quadrant. Then the value of  $3\alpha$  is equal to \_\_\_\_\_.

**Answer (22)**

**Sol.**



$$A_1 = \int_0^2 2\sqrt{2}\sqrt{x} - x dx$$

$$= 2\sqrt{2} \times \frac{2}{3} \cdot x^{\frac{3}{2}} - \frac{x^2}{2} \Big|_0^2$$

$$= \frac{4\sqrt{2}}{3} \times 2\sqrt{2} - 2$$

$$= \frac{16}{3} - 2 = \frac{10}{3}$$

$$A_2 = \int_2^8 2\sqrt{2}\sqrt{x} - x dx = \frac{4\sqrt{2}}{3} \cdot x^{\frac{3}{2}} - \frac{x^2}{2} \Big|_2^8$$

$$= \frac{4\sqrt{2}}{3} (16\sqrt{2} - 2\sqrt{2}) - 30$$

$$= \frac{112}{3} - 30 = \frac{22}{3}$$

$$A_2 > A_1 \Rightarrow 3\alpha = 22$$

